

Original Article

Biomarkers and Choice of Biologic Agent: A Comparison of the Efficacy of Tocilizumab and Etanercept in Rheumatoid Arthritis Patients Presenting with Leukocytosis or Thrombocytosis

Kiyomitsu Miyachi ^{1,2}, Belinda Sasse ^{1,3}, So Nomoto ⁴

¹Department of Rheumatology, Keigu Clinic, 2-2 Ichibanishinaka-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0023, Japan

²First Diagnostic Division, Health Sciences Research Institute, Yokohama, Kanagawa 240-0005, Japan

³Department of Rheumatology, Monash Health, 246 Clayton Rd, Clayton, Victoria 3168, Australia

⁴Department of Orthopedic Surgery, Saiseikai Yokohama City Eastern Hospital, Yokohama, Kanagawa, 230-0012, Japan

*Corresponding author: Dr. Kiyomitsu Miyachi, M.D. Keigu Clinic, 2-2 Ichibanishinaka-cho Tsurumi-ku Yokohama, Kanagawa 230-0023, Japan, E-mail: mkiyomitsumd_8@hotmail.com, Tel: +81 45 501 5361


Received: 03-31-2015

Accepted: 09-05-2015

Published:

Copyright: © 2015 Kiyomitsu


RA Cytokine Network



Nishimoto, N: Riumachika, 35(5):459-463, 2006.



Cytokine; IL-6 function



Nishimoto N et al. Igakunoayumi 188;942-948:1999 revised